arXiv:1604.02489v1 [math.DS] 8 Apr 2016

$\mathbf{IP}_\mathbf{r}^*$ recurrence and nilsystems

V. Bergelson and A. Leibman

April 12, 2016

Abstract

We prove that nilsystems possess the IP_r^*-recurrence property and, moreover, are characterized by this property.

0. Introduction

Our goal in this paper is to provide, in terms of recurrence, a characteristic of *nilsystems*, namely, systems of the form (X,T) where X is a *nilmanifold* (that is, a compact homogeneous space of a nilpotent Lie group G) and T is a *niltranslation* (that is, a translation of X defined by an element of G). The motivation for this study comes from the fact that nilsystems are intrincically related to various problems arising in ergodic theory of multiple recurrence, combinatorics, and number theory, and understanding the recurrence properties of niltranslations leads to interesting applications in these areas.

A system (X,T), where X is a compact metric space and T is a self-homeomorphism of X, is said to be *distal* if for any distinct $x,y\in X$, $\inf_{n\in\mathbb{Z}}\operatorname{dist}(T^nx,T^ny)>0$. Given a sequence $n_1,n_2,\dots$ in $\mathbb{Z}$, the set $\{n_{i_1}+\cdots+n_{i_s}: s\in\mathbb{N},\ i_1<\cdots<i_s\}$ of finite sums of distinct elements of this sequence is called an *IP-set*. A subset E of $\mathbb{Z}$ is called an *IP*-set* if it has a nonempty intersection with every IP-set. H. Furstenberg had shown that distal systems are characterized by *the IP*-recurrence property*:

Theorem 0.1. ([F], Theorem 9.11) *A system (X,T) is distal iff for any $x_0\in X$ and any neighborhood U of x_0 the set of returns $R_U(x_0)=\{n\in\mathbb{Z}: T^nx_0\in U\}$ of x_0 to U is an IP*-set.*

It is well known that nilsystems are distal (see [AGH], [Ke1], [Ke2]), and thus are IP*-recurrent. A special class of nilsystems, the affine skew product transformations on tori[(1)] have a stronger recurrence property, namely, that of *IP_0^*-recurrence*. For an integer $r\in\mathbb{N}$ and an r-element sequence $n_1,\dots,n_r$ in $\mathbb{Z}$, we call the set $\{n_{i_1}+\cdots+n_{i_s}: 1\le s\le r,\ i_1<\cdots<i_s\}$ of sums of distinct elements of this sequence an *IP_r-set*. A set $E\subseteq\mathbb{Z}$ is called an *IP_r^*-set* if it has a nonempty intersection with every IP_r-set in $\mathbb{Z}$. We say that a set is an

Partially supported by NSF grants DMS-1162073 and DMS-1500575.

(1) An affine skew product transformation of the k-dimensional torus $\mathbb{T}^k=\mathbb{R}^k/\mathbb{Z}^k$ is defined by the formula $T(x_1,\dots,x_k)=(x_1+\alpha_1,x_2+a_{2,1}x_1+\alpha_2,\dots,x_k+a_{k,k-1}x_{k-1}+\cdots+a_{k,1}x_1+\alpha_k)$ with $\alpha_i\in\mathbb{T}$ and $a_{i,j}\in\mathbb{Z}$.

IP_0^*-set if it is an IP_r^* set for some $r \in \mathbb{N}$. It follows from [B], Theorem 7.7, that any affine skew product transformation of a torus has the IP_0^*-recurrence property: for every $x_0 \in \mathbb{T}^k$ and any open neighborhood U of x_0 the set $R_U(x_0) = \{n \in \mathbb{Z} : T^n x_0 \in U\}$ is an IP_0^*-set. One can show that not every distal system is IP_0^*-recurrent (see [BL3], Section 1), and it is natural to ask what distal systems are characterized by the property of IP^*-recurrence. Let us say that a system (X, T) is *a pre-nilsystem* if it is an inverse limit of a sequence of nilsystems. (Notice that, in contrast with the definition of the so-called *pro-nilsystems*, in the definition of pre-nilsystems we don't require the nilpotency class of the nilsystems be bounded.) We prove:

Theorem 0.2. *Any pre-nilsystem is IP_0^*-recurrent. Any IP_0^*-recurrent system is a disjoint union of pre-nilsystems.*

The second statement of the theorem is an easy corollary of the results from [HKM]. To prove the first statement, we use a coordinate approach. On any nilmanifold X one has natural coordinates such that under the action of a niltranslation T the sequence of coordinates of the image $T^n x_0$ of any point $x_0 \in X$ is given by *generalized polynomials* (see [BL2], Theorem A). We therefore need to deal with images of IP-sets under generalized polynomial mappings, which is a special case of *generalized polynomial IP-sets*. Conventional IP- and IP_r-sets in $\mathbb{Z}$ can be viewed as the images of mappings $\varphi\colon \mathcal{F}(A) \longrightarrow \mathbb{Z}$ from the semigroup $\mathcal{F}(A)$ of finite subsets of A, for $A = \mathbb{N}$ and, respectively, for $A = \{1, \dots, r\}$, defined by $\varphi(\alpha) = \sum_{i\in\alpha} a_i$. Such a mapping φ is "linear" in the followig sense: $\varphi(\alpha \cup \beta) = \varphi(\alpha) + \varphi(\beta)$ whenever $\alpha, \beta \in \mathcal{F}(A)$ are disjoint. Let H be an additive abelian group; one may introduce the notion of polynomial mappings $\mathcal{F}(A) \longrightarrow H$ as follows. For a mapping $\varphi\colon \mathcal{F}(A) \longrightarrow H$ and a set $\beta \in \mathcal{F}(A)$ let the "β-derivative" $D_\beta\varphi$ be the mapping $\mathcal{F}(A \setminus \beta) \longrightarrow H$ defined by $D_\beta\varphi(\alpha) = \varphi(\alpha + \beta) - \varphi(\alpha)$. Then we say that a mapping $\varphi\colon \mathcal{F}(\{1, \dots, r\}) \longrightarrow H$ is *polynomial of degree* $\le d$ if for any disjoint $\beta_0, \beta_1, \dots, \beta_d \in \mathcal{F}(\{1, \dots, r\})$, $D_{\beta_0} D_{\beta_1} \cdots D_{\beta_d}\varphi = 0$. (See [BL1], Section 8.1.) Examples of quadratic (that is, of degree ≤ 2) polynomial mappings are $\varphi(\alpha) = \big(\sum_{i\in\alpha} a_i\big)^2$, $\varphi(\alpha) = \big(\sum_{i\in\alpha} a_i\big)\big(\sum_{i\in\alpha} b_i\big) = \sum_{i,j\in\alpha} a_i b_j$, and $\varphi(\alpha) = \sum_{i,j\in\alpha} c_{i,j}$, where $a_i, b_j, c_{i,j} \in H$. *Generalized polynomial mappings* are the mappings "made" of (conventional) polynomial mappings using the operations of addition, multiplication, and taking the integer part. (An example is $\varphi = \big[[\varphi_1]\varphi_2 + \varphi_3\big]\varphi_4 + [\varphi_5][\varphi_6]\varphi_7$, which is "made" of the polynomial mappings $\varphi_1, \dots, \varphi_7$.) Let us define *the total degree* of a generalized polynomial mapping φ as the sum $D = \sum_i \deg\varphi_i$ of the degrees of all the "conventional" polynomial mappings φ_i of which φ is made, and let us say that a generalized polynomial mapping is *constant free* if all the φ_i "vanish at $\emptyset$": $\varphi_i(\emptyset) = 0$. For $x \in \mathbb{R}$ let $\|x\| = \mathrm{dist}(x, \mathbb{Z})$. We prove:

Theorem 0.3. (Cf. Theorem 1.11 below.) *For any $D \in \mathbb{N}$ and $\varepsilon > 0$ there exists $r = r(D, \varepsilon) \in \mathbb{N}$ such that for any constant-free generalized polynomial mapping $\varphi\colon \{1, \dots, r\} \longrightarrow \mathbb{R}$ of total degree $\le D$ there exists a nonempty $\alpha \subseteq \{1, \dots, r\}$ for which $\|\varphi(\alpha)\| < \varepsilon$.*

The *VIP-sets* in $\mathbb{Z}^l$ are defined as the images $\{\varphi(\alpha) : \alpha \in \mathcal{F}(\mathbb{N}),\ \alpha \ne \emptyset\}$ of polynomial mappings $\varphi\colon \mathcal{F}(\mathbb{N}) \longrightarrow \mathbb{Z}^l$ with $\varphi(\emptyset) = 0$, and we say that a set $E \subseteq \mathbb{Z}^l$ is *a VIP^*-set* if E has a nonempty intersection with every VIP-set in $\mathbb{Z}^l$. Similarly, for all $d, r \in \mathbb{N}$, we

define *VIP$_{d,r}$-sets* as the images $\{\varphi(\alpha) : \alpha \subseteq \{1,\dots,r\},\ \alpha \neq \emptyset\}$ of polynomial mappings $\varphi\colon \mathcal{F}(\{1,\dots,r\}) \longrightarrow \mathbb{Z}^l$ of degree $\leq d$ and with $\varphi(\emptyset) = 0$, and say that a set $E \in \mathbb{Z}^l$ is a *VIP$^*_{d,r}$-set* if it has a nonempty intersection with every VIP$_{d,r}$-set. We will also say that a set $E \in \mathbb{Z}^l$ is a *VIP$^*_0$-set* if for any $d \in \mathbb{N}$, E is an VIP$^*_{d,r}$-set for some $r \in \mathbb{N}$. Theorem 0.3 now implies the following result:

Theorem 0.4. *For any $D, d \in \mathbb{N}$ and $\varepsilon > 0$ there exists $r = r(D,d,\varepsilon) \in \mathbb{N}$ such that for any $l \in \mathbb{N}$ and any constant-free generalized polynomial mapping $\varphi\colon \mathbb{Z}^l \longrightarrow \mathbb{R}$ of total degree $\leq D$ the set $\{n \in \mathbb{Z}^l : \|\varphi(n)\| < \varepsilon\}$ is a VIP$^*_{d,r}$-set.*

Let G be a nilpotent Lie group; an *l-parameter polynomial sequence* in G is a mapping $g\colon \mathbb{Z}^l \longrightarrow G$ of the form $T_1^{p_1(n)} \cdots T_b^{p_b(n)}$, $n \in \mathbb{Z}^l$, where $T_i \in G$ and p_i are polynomials $\mathbb{Z}^l \longrightarrow \mathbb{Z}$; the *naive degree* of g is defined as $\max_i \deg p_i$. Using the fact that the coordinates of a point of a nilmanifold under the action of a polynomial sequence of niltranslations are generalized polynomials, we obtain as a corollary of Theorem 0.4 the following strengthening of the first part of Theorem 0.2:

Theorem 0.5. (Cf. Theorem 1.12 below.) *Let X be a nilmanifold with metric ρ (compatible with the homogeneous space structure on X). For any $a, d \in \mathbb{N}$ and $\varepsilon > 0$ there exists $r = r(a,d,\varepsilon) \in \mathbb{N}$ such that for any $x_0 \in X$, any $l \in \mathbb{N}$, and any l-parameter polynomial sequence g of niltranslations of X of naive degree $\leq a$ and with $g(0) = \mathrm{Id}_X$, the set $R_U(x_0) = \{n \in \mathbb{Z}^l : \rho(g(n)x_0, x_0) < \varepsilon\}$ is a VIP$^*_{d,r}$-set.*

We say that a dynamical system (X,T) is *VIP*-recurrent* if for any $x_0 \in X$ and any neighborhood U of x_0 the set of returns $R_U(x_0) = \{n \in \mathbb{Z} : T^n x_0 \in U\}$ is a VIP*-set, and is *VIP$^*_0$-recurrent* if for any $x_0 \in X$ and any neighborhood U of x_0 the set $R_U(x_0)$ is a VIP$^*_0$-set. The VIP*-recurrence property turns out to be strictly stronger than that of the IP*-recurrence: there exist distal but not VIP*-recurrent systems.[(2)] As for the VIP$^*_0$-recurrence, the assertion of Theorem 0.5 implies that pre-nilsystems possess this property, and so by Theorem 0.2, the VIP$^*_0$-recurrence is equivalent to the IP$^*_0$-recurrence.

1. VIP$^*_0$-recurrence of bounded generalized polynomials

Let A be a set and $(H,+)$ be an abelian group. For $r \in \mathbb{N}$ we will denote by $[1,r]$ the interval $\{1,\dots,r\}$ in $\mathbb{N}$. We denote by $\mathcal{F}(A)$ the set of finite subsets of A, by $A^{(d)}$, $d \in \mathbb{N}$, the set of subsets of A of cardinality d, and by $A^{(\leq d)}$, $d \in \mathbb{N}$, the set of nonempty subsets of A of cardinality $\leq d$, $A^{(\leq d)} = \bigcup_{l=1}^d A^{(l)}$.

We start with defining polynomial mappings on $\mathcal{F}(A)$. We say that a mapping $\varphi\colon \mathcal{F}(A) \longrightarrow H$ is *linear* if it satisfies the identity $\varphi(\alpha \cup \beta) = \varphi(\alpha) + \varphi(\beta)$ whenever $\alpha, \beta \in \mathcal{F}(A)$ are disjoint, and will denote the set of linear mappings $\mathcal{F}(A) \longrightarrow H$ by $\mathrm{Lin}(A,H)$. A mapping $\varphi \in \mathrm{Lin}(A,H)$ is uniquely defined by its values at singletons: for any $\alpha \in \mathcal{F}(A)$, $\varphi_\alpha = \sum_{a\in\alpha} \widehat{\varphi}(\{a\})$. We will call the mapping $\widehat{\varphi}\colon A \longrightarrow H$ defined by $\widehat{\varphi}(a) = \varphi(\{a\})$ *the producing function* for φ; we then have $\varphi(\alpha) = \sum_{a\in\alpha} \widehat{\varphi}(a)$, $\alpha \in \mathcal{F}(A)$.

(2) See [P], Corollary 5.1, where it is shown that for any nonlinear polynomial $p\colon \mathbb{Z} \longrightarrow \mathbb{Z}$ there exists an affine skew product transformation T such that $\liminf_n \mathrm{dist}(T^{p(n)}0, 0) > 0$.

For a mapping $\varphi\colon \mathcal{F}(A) \longrightarrow H$ and $\beta \in \mathcal{F}(A)$ we define *the β-derivative* $D_\beta\varphi$ of φ by $D_\beta\varphi(\alpha) = \varphi(\alpha\cup\beta) - \varphi(\alpha)$, $\alpha \in \mathcal{F}(A\setminus\beta)$. We say that a mapping φ is *polynomial of degree* $\leq d$ if for any $d+1$ pairwise disjoint sets $\beta_0, \ldots, \beta_d \in \mathcal{F}(A)$ one has $D_{\beta_d}\cdots D_{\beta_0}\varphi = 0$.

We will denote by $\mathrm{Pol}_d(A, H)$ the group of polynomial mappings $\mathcal{F}(A) \longrightarrow H$ of degree $\leq d$. We will mainly deal with polynomial mappings "having zero constant term"; let us denote by $\mathrm{Pol}^0_d(A, H)$ the subgroup $\left\{\varphi \in \mathrm{Pol}_d(A, H) : \varphi(\emptyset) = 0\right\}$ of $\mathrm{Pol}_d(A, H)$. Notice that $\mathrm{Lin}(A, H) = \mathrm{Pol}^0_1(A, H)$.

One can show (see [BL1], sections 8.3-8.5) that any polynomial mapping $\varphi \in \mathrm{Pol}^0_d(A, H)$ can be represented in the form $\varphi(\alpha) = \Phi(\alpha^d)$, $\alpha \in \mathcal{F}(A)$, for some mapping $\Phi \in \mathrm{Lin}(A^d, H)$, so that

$$\varphi(\alpha) = \sum_{v\in\alpha^d} \widehat{\Phi}(v),\ \alpha \in \mathcal{F}(A), \tag{1.1}$$

where $\widehat{\Phi}\colon A^d \longrightarrow H$ is the producing function for Φ. We will call $\widehat{\Phi}$ *a q-producing function for* φ.

The q-producing function for a polynomial mapping $\varphi \in \mathrm{Pol}^0_d(A, H)$ is not canonically defined. A more natural is *the t-producing function* for φ, a function $\widetilde{\Phi}\colon A^{(\leq d)} \longrightarrow H$ such that for any $\alpha \in \mathcal{F}(A)$,

$$\varphi(\alpha) = \sum_{u\in\alpha^{(\leq d)}} \widetilde{\Phi}(u).$$

The t-producing function $\widetilde{\Phi}$ for φ is defined uniquely (and provides a natural definition of polynomial mappings in the case H is a commutative semigroup). In terms of $\widetilde{\Phi}$, φ is the sum of its *homogeneous components*, $\varphi = \varphi_1 + \cdots + \varphi_d$, where for each i, $\varphi_i(\alpha) = \sum_{\delta\in\alpha^{(d)}} \widetilde{\Phi}(\delta)$. To obtain the t-producing function $\widetilde{\Phi}$ from a q-producing function $\widehat{\Phi}$ for φ one simply sums up the values of $\widehat{\Phi}$ at the elements of $A^{(d)}$ corresponding to the same element of $A^{(\leq d)}$: for any $u \in A^{(\leq d)}$,

$$\widetilde{\Phi}(u) = \sum_{\substack{v=(a_1,\ldots,a_d)\in\alpha^d \\ \{a_1,\ldots,a_d\}=u}} \widehat{\Phi}(v). \tag{1.2}$$

Let B be a collection of pairwise disjoint finite subsets of A; we will call B *a disjoint subcollection* of A; if $|B| = s$ we will say that B is *a disjoint s-subcollection.* Given a disjoint subcollection B of A, we have an injection $\mathcal{F}(B) \longrightarrow \mathcal{F}(A)$ defined by $\gamma \mapsto \bigcup\gamma$, and we will identify $\mathcal{F}(B)$ with its image in $\mathcal{F}(A)$. Given a polynomial mapping $\varphi\colon \mathcal{F}(A) \longrightarrow H$, we call the polynomial mapping $\varphi|_{\mathcal{F}(B)}$ *a subpolynomial* of φ corresponding to the disjoint subcollection B and denote it by $\varphi_{\downarrow B}$. Any disjoint subcollection B of a disjoint subcollection of A induces the disjoint subcollection $B' = \left\{\bigcup C : C \in B\right\}$ of A; abusing notation, we will denote the subpolynomial $\varphi_{\downarrow B'}$ of φ by $\varphi_{\downarrow B}$.

Let $\widehat{\Phi}\colon A^d \longrightarrow H$ be a q-producing function for a polynomial mapping $\varphi\colon \mathcal{F}(A) \longrightarrow H$ of degree $\leq d$ and let $\Phi \in \mathrm{Lin}(A^d, H)$ be the linear mapping produced by $\widehat{\Phi}$. Given a disjoint s-subcollection $B = \{B_1, \ldots, B_s\}$ of A, one finds a q-producing function for the

subpolynomial $\varphi_{\downarrow B}$ as follows. For any $\beta \subseteq B$ we have

$$\begin{aligned}\varphi_{\downarrow B}(\beta) = \varphi\Big(\bigcup_{C\in\beta} C\Big) = \Phi\Big(\Big(\bigcup_{C\in\beta} C\Big)^d\Big) = \sum_{v\in(\bigcup_{C\in\beta} C)^d} \widehat{\Phi}(v) \\ = \sum_{C_1,\dots,C_d\in\beta} \sum_{v\in C_1\times\cdots\times C_d} \widehat{\Phi}(v) = \sum_{(C_1,\dots,C_d)\in\beta^d} \Phi(C_1,\dots,C_d);\end{aligned} \tag{1.3}$$

thus, the mapping $\Phi_{|B^d}$ is a q-producing function for $\varphi_{\downarrow B}$.

The following proposition establishes the IP_r^*-recurrence property of polynomial mappings with values in the torus $\mathbb{T} = \mathbb{R}/\mathbb{Z}$:

Proposition 1.1. (Cf. [B], Theorem 7.7) *For any $k, d \in \mathbb{N}$ and $\varepsilon > 0$ there exists $r = r(k, d, \varepsilon) \in \mathbb{N}$ such that for any $\varphi_1, \dots, \varphi_k \in \mathrm{Pol}_d^0([1, r], \mathbb{T})$ there exists a nonempty $\alpha \in \mathcal{F}([1, r])$ such that $\mathrm{dist}(\varphi_i(\alpha), 0) < \varepsilon$ for all $i \in \{1, \dots, k\}$.*

Proof. Put $c = \lceil 1/\varepsilon \rceil$ and partition the torus $\mathbb{T}$ into c intervals of length $\leq 1/\varepsilon$. By the Polynomial Hales-Jewett theorem (see [BL1], Theorem 0.10), there exists $r \in \mathbb{N}$ such that for any partition of $\mathcal{F}([1, r]^d \times [k])$ into c subsets there exist $\gamma \subset [1, r]^d \times [k]$ and a nonempty $\alpha \subseteq [1, r]$ such that $\gamma \cap (\alpha^d \times [k]) = \emptyset$ and the sets $\gamma, \gamma \cup (\alpha^d \times \{1\}), \dots, \gamma \cup (\alpha^d \times \{k\})$ belong to the same element of the partition. Let $\varphi_1, \dots, \varphi_k \in \mathrm{Pol}_d^0([1, r], \mathbb{T})$. For each i let $\widehat{\Phi}_i \colon [1, r]^d \longrightarrow \mathbb{T}$ be a q-producing function for φ_i. Define a mapping $\widehat{\Phi} \colon [1, r]^d \times [k] \longrightarrow \mathbb{T}^k$ by $\widehat{\Phi}(v, i) = \widehat{\Phi}_i(v)$, $v \in [1, r]^d$, $i \in [k]$, and let $\Phi \in \mathrm{Lin}([1, r]^d \times [k], \mathbb{T})$ be the linear mapping produced by $\widehat{\Phi}$. Then, via Φ, the partition of $\mathbb{T}$ defines a partition of $\mathcal{F}([1, r]^d \times [k])$ into c subsets. Applying the Polynomial Hales-Jewett theorem, we can find $\gamma \subset [1, r]^d \times [k]$ and a nonempty $\alpha \subseteq [1, r]$ such that $\gamma \cap (\alpha^d \times [k]) = \emptyset$ and the sets $\gamma, \gamma \cup (\alpha^d \times \{1\}), \dots, \gamma \cup (\alpha^d \times \{k\})$ belong to the same element of the partition; then for any i, $\Phi(\gamma)$ and $\Phi(\gamma \cup (\alpha^d \times \{i\}))$ belong to the same partition of $\mathbb{T}$, and so, $\mathrm{dist}\big(\Phi(\gamma), \Phi(\gamma \cup (\alpha^d \times \{i\}))\big) < \varepsilon$. Since $\Phi(\gamma \cup (\alpha^d \times \{i\})) = \Phi(\gamma) + \Phi(\alpha^d \times \{i\}) = \Phi(\gamma) + \varphi_i(\alpha)$, this implies that $\mathrm{dist}(0, \varphi_i(\alpha)) < \varepsilon$. ■

Let $\|x\|$ denote the distance from $x \in \mathbb{R}$ to $\mathbb{Z}$. We may then reformulate Proposition 1.1 as follows:

Corollary 1.2. *For any $k, d \in \mathbb{N}$ and $\varepsilon > 0$ there exists $r = r(k, d, \varepsilon) \in \mathbb{N}$ such that for any $\varphi_1, \dots, \varphi_k \in \mathrm{Pol}_d^0([1, r], \mathbb{R})$ there exists a nonempty $\alpha \in \mathcal{F}([1, r])$ such that $\|\varphi_i(\alpha)\| < \varepsilon$ for all $i \in \{1, \dots, k\}$.*

Next we show that if r is large enough, any polynomial mapping $\varphi \in \mathrm{Pol}_d^0([1, r], \mathbb{T})$ has a subpolynomial whose q-producing function is arbitrarily small:

Proposition 1.3. *For any $d, s \in \mathbb{N}$ and $\varepsilon > 0$ there exists $r \in \mathbb{N}$ such that for any $\varphi \in \mathrm{Pol}_d^0([1, r], \mathbb{T})$ there exists a disjoint s-subcollection B of $[1, r]$ such that a q-producing function $\widehat{\Phi}_B$ for $\varphi_{\downarrow B}$ satisfies $\mathrm{dist}(\widehat{\Phi}_B, 0) < \varepsilon$.*

Proof. Take $r_0 = r(s^d, d, \varepsilon)$ as in Corollary 1.2, and put $A = [1, s] \times [1, r_0]$ (and $r = |A| = sr_0$). Let $\varphi \in \mathrm{Pol}_d^0([1, r], \mathbb{T})$. Let $\widehat{\Phi}\colon [1, r] \longrightarrow \mathbb{T}$ be a q-producing function for φ and let $\Phi \in \mathrm{Lin}([1, r]^d, \mathbb{T})$ be the linear mapping produced by $\widehat{\Phi}$. For each $i = (i_1, \dots, i_d) \in [1, s]^d$ define a polynomial mapping $\varphi_I \in \mathrm{Pol}_d^0([1, r_0], \mathbb{T})$ by $\varphi_I(\alpha) = \Phi((\{i_1\}\times\alpha)\times\cdots\times(\{i_d\times\alpha)\})$. By Corollary 1.2 there exists $\alpha \subseteq [1, r_0]$ such that $\mathrm{dist}(\varphi_I(\alpha), 0) < \varepsilon$ for all $I \in [1, s]^d$. Take the disjoint s-subcollection $B = \big\{\{i\} \times \alpha : i \in [1, s]\big\}$ of A. By the choice of α, for any $w \in B^d$ we have $\mathrm{dist}(\Phi(w), 0) < \varepsilon$. Since, by (1.3), $\Phi_{|_{B^d}}$ is a q-producing function for $\varphi_{\downarrow_B}$, we are done. ■

Replacing in Proposition 1.3 ε by ε/s^d, we obtain:

Corollary 1.4. *For any $d, s \in \mathbb{N}$ and $\varepsilon > 0$ there exists $r \in \mathbb{N}$ such that for any $\varphi \in \mathrm{Pol}_d^0([1, r], \mathbb{T})$ there exists a disjoint s-subcollection B of $[1, r]$ such that $\mathrm{dist}(\varphi_{\downarrow_B}, 0) < \varepsilon$.*

By formula (1.2), any value of the t-producing function for $\varphi \in \mathrm{Pol}_d^0(A, \mathbb{R})$ is a sum of less than d^d values of the q-producing function for φ. Hence, Proposition 1.3 implies the following its version:

Proposition 1.5. *For any $d, s \in \mathbb{N}$ and $\varepsilon > 0$ there exists $r \in \mathbb{N}$ such that for any $\varphi \in \mathrm{Pol}_d^0([1, r], \mathbb{T})$ there exists a disjoint s-subcollection B of $[1, r]$ such that the t-producing function $\widetilde{\Phi}_B$ for $\varphi_{\downarrow_B}$ satisfies $\mathrm{dist}(\widetilde{\Phi}_B, 0) < \varepsilon$.*

In terms of polynomial mappings with values in $\mathbb{R}$, Proposition 1.5 takes the following form:

Corollary 1.6. *For any $d, s \in \mathbb{N}$ and $\varepsilon > 0$ there exists $r \in \mathbb{N}$ such that for any $\varphi \in \mathrm{Pol}_d^0([1, r], \mathbb{R})$ there exists a disjoint s-subcollection B of $[1, r]$ such that the t-producing function $\widetilde{\Phi}_B$ for $\varphi_{\downarrow_B}$ satisfies $\|\widetilde{\Phi}_B\| < \varepsilon$.*

Now let $\varphi \in \mathrm{Pol}_d^0([1, r], \mathbb{R})$ be a polynomial mapping whose t-producing function $\widetilde{\Phi}$ satisfies $\|\widetilde{\Phi}\| < 1/r^d$. We will denote by $[x]$ the integer and by $\{x\}$ the fractional parts of $x \in \mathbb{R}$. If $x \in \mathbb{R}$ satisfies $\|x\| < \varepsilon$, then either $\{x\} < \varepsilon$ or $\{x\} > 1 - \varepsilon$. If $x_1, \dots, x_n \in \mathbb{R}$ satisfy $\{x_i\} < 1/n$, $i = 1, \dots, n$, then $\big[\sum_{i=1}^n x_i\big] = \sum_{i=1}^n [x_i]$. Thus, if $\widetilde{\Phi}$ satisfies $\{\widetilde{\Phi}\} < 1/r^d$, then for any $\alpha \subseteq [1, r]$,

$$[\varphi(\alpha)] = \Big[\sum_{u\in\alpha^{(\le d)}} \widetilde{\Phi}(u)\Big] = \sum_{u\in\alpha^{(\le d)}} [\widetilde{\Phi}(u)]$$

and so, $[\varphi]$ is also a polynomial mapping, $[\varphi] \in \mathrm{Pol}_d^0([1, r], \mathbb{Z})$, with the t-producing function $[\widetilde{\Phi}]$.

For any $x \in \mathbb{R} \setminus \mathbb{Z}$, $[x] = -[-x] - 1$ and $\{-x\} = 1 - \{x\}$, so, if $x_1, \dots, x_n \in \mathbb{R}$ satisfy $\{x_i\} > 1 - 1/n$, $i = 1, \dots, n$, then

$$\big[\textstyle\sum_{i=1}^n x_i\big] = -\big[-\sum_{i=1}^n x_i\big] - 1 = -\big[\sum_{i=1}^n (-x_i)\big] - 1 = -\sum_{i=1}^n [-x_i] - 1 = \sum_{i=1}^n (-[-x_i]) - 1.$$

Applying this to $\widetilde{\Phi}$, we see that if $\widetilde{\Phi}$ satisfies $\{\widetilde{\Phi}\} > 1 - 1/r^d$, then for any $\alpha \subseteq [1, r]$,

$$[\varphi(\alpha)] = \Big[\sum_{u\in\alpha^{(\le d)}} \widetilde{\Phi}(u)\Big] = \sum_{u\in\alpha^{(\le d)}} \Big(-[-\widetilde{\Phi}(u)]\Big) - 1.$$

So, $[\varphi]+1$ is a polynomial mapping, $[\varphi]+1 \in \mathrm{Pol}_d^0([1,r],\mathbb{Z})$, with the t-producing function $-[-\widetilde{\Phi}]$.

In the general case, when $\|\widetilde{\Phi}\| < 1/r^d$, we may have neither $\{\widetilde{\Phi}\} < 1/r^d$ nor $\{\widetilde{\Phi}\} > 1 - 1/r^d$. However, if φ is a homogeneous polynomial of degree $l \le d$ (which means that $\varphi(\alpha) = \sum_{u\in\alpha^{(l)}} \widetilde{\Phi}(u)$), then, given $s \in \mathbb{N}$, if r is large enough, by the classical Ramsey theorem we can choose an s-element subset B of $[1,r]$ such that either $\{\widetilde{\Phi}(u)\} < 1/r^d$ for all $u \in B^{(d)}$ or $\{\widetilde{\Phi}(u)\} > 1 - 1/r^d$ for all $u \in B^{(d)}$. Identifying B with the "singleton disjoint '-subcollection' $\{\{b\} : b \in B\}$ of $[1,r]$, we will therefore have $[\varphi_{\downarrow B}] \in \mathrm{Pol}_d^0(B,\mathbb{Z}) + e$ with $e \in \{0,-1\}$.

For a general $\varphi \in \mathrm{Pol}_d^0([1,r],\mathbb{R})$, applying this argument to all homogeneous components of φ and using a diagonal process, we arrive at the following lemma:

Lemma 1.7. *For any $d, s \in \mathbb{N}$ there exists $r \in \mathbb{N}$ such that for any $\varphi \in \mathrm{Pol}_d^0([1,r],\mathbb{R})$ whose t-producing function $\widetilde{\Phi}$ satisfies $\|\widetilde{\Phi}\| < 1/r^d$ there exists a (singleton) disjoint subcollection B of $[1,r]$ such that $[\varphi_{\downarrow B}] \in \mathrm{Pol}_d^0(B,\mathbb{Z}) + e$ with $e \in \{0,-1,\dots,-d\}$.*

Combining Lemma 1.7 with Corollary 1.6 we obtain:

Theorem 1.8. *For any $d, s \in \mathbb{N}$ there exists $r \in \mathbb{N}$ such that for any $\varphi \in \mathrm{Pol}_d^0([1,r],\mathbb{R})$ there exists a disjoint s-subcollection B of $[1,r]$ such that $[\varphi] \in \mathrm{Pol}_d^0(B,\mathbb{Z}) + e$ with $e \in \{0,-1,\dots,-d\}$.*

Using induction on k, one can extend Theorem 1.8 to the case of k polynomials:

Theorem 1.9. *For any $k, d_1, \dots, d_k, s \in \mathbb{N}$ there exists $r = r(k,(d_1,\dots,d_k),s) \in \mathbb{N}$ such that for any $\varphi_i \in \mathrm{Pol}_{d_i}^0([1,r],\mathbb{R})$, $i = 1,\dots,k$, there exists a disjoint s-subcollection B of $[1,r]$ such that for every $i \in \{1,\dots,k\}$, $[\varphi_i] \in \mathrm{Pol}_{d_i}^0(B,\mathbb{Z}) + e_i$, with $e_i \in \{0,-1,\dots,-d\}$.*

A generalized polynomial is a function obtained from conventional polynomials using the operations of taking the integer part, addition, and multiplication. We say that a generalized polynomial φ is *constant free* if all polynomials involved in the expression of φ have zero constant term. (More exactly, a generalized polynomial is constant free if it has *a representation* in which all polynomials have zero constant term. A similar convention applies to all the definitions below.) We say that a polynomial φ is *open* if it has no summand that is a product of "closed" generalized polynomials $[\varphi_i]$; formally, φ is open if it is a conventional polynomial, or $\varphi = \varphi_1 + \varphi_2$ where φ_1 and φ_2 are open, or $\varphi = \varphi_1\varphi_2$ where at least one of φ_1 and φ_2 is open. Under these definitions, any open constant-free generalized polynomial is representable in the form

$$\varphi = \sum_{j=1}^{m} [\varphi_{j,1}] \cdots [\varphi_{j,l_j}]\varphi_{j,0} \tag{1.4}$$

where for every j, $\varphi_{j,1},\dots,\varphi_{j,l_j}$ are open constant-free generalized polynomials and $\varphi_{j,0}$ are conventional polynomials with zero constant term.

We now introduce the notions of height, width, and degree for a generalized polynomial φ:

The height $h(\varphi)$ of φ is the maximum length of sequences of nested brackets in φ: we put $h(\varphi) = 0$ if φ is a conventional polynomial and we say that $h(\varphi) \leq h$ if φ has a representation (1.4) where for all j and all $t \geq 1$, $h(\varphi_{j,t}) \leq h-1$.
The width $w(\varphi)$ is the maximum number of components in φ itself and in all its components: we put $w(\varphi) = 1$ if φ is a conventional polynomial and we say that $w(\varphi) \leq w$ if φ has a representation (1.4) where $w(\varphi_{j,t}) \leq w$ for all j and all $t \geq 1$ and also $\sum_{j=1}^{m}(l_j+1) \leq w$.
The degree $d(\varphi)$ of φ is defined as usual under the assumption that $\deg[\varphi] = \deg\varphi$: we say that $d(\varphi) \leq d$ if φ has a representation (1.4) with $\max_{j=1}^{m}\left(\sum_{t=0}^{l_j}\deg\varphi_{j,t}\right) \leq d$.
(For example, for $\varphi(x) = [[x^2+1]x][x^3+2x]x+[x^2](x+1)+x^3$ we have $h(\varphi) = 2$, $w(\varphi) = 6$, and $d(\varphi) = 7$.)

We extend the above definitions to generalized polynomial mappings with domain $\mathcal{F}(A)$, and will denote by $\mathrm{GPol}^0_{d,h,w}(A,H)$ the set (the algebra) of open constant-free generalized polynomial mappings $\varphi\colon \mathcal{F}(A) \longrightarrow H$, where $H = \mathbb{R}$ or $\mathbb{Z}$, with $d(\varphi) \leq d$, $h(\varphi) \leq h$, and $w(\varphi) \leq w$. Given $\varphi \in \mathrm{GPol}^0_{d,h,w}(A,H)$ and a disjoint subcollection B of A, we define the generalized polynomial mapping $\varphi_{\downarrow B} \in \mathrm{GPol}^0_{d,h,w}(B,H)$ as the restriction of φ to the set $\mathcal{F}(B)$ considered as a subset of $\mathcal{F}(A)$.

The following theorem says that generalized polynomial mappings turn into ordinary polynomial mappings after being restricted to a suitable disjoint subcollection of their domain:

Theorem 1.10. *For any* $k, d_1, \ldots, d_k, h, w, s \in \mathbb{N}$ *there exists* $r = r(k,(d_1,\ldots,d_k),h,w,s) \in \mathbb{N}$ *such that for any* $\varphi_i \in \mathrm{GPol}^0_{d_i,h,w}([1,r],\mathbb{R})$, $i = 1,\ldots,k$, *there exists a disjoint* s*-subcollection* B *of* $[1,r]$ *such that* $\varphi_{i\downarrow B} \in \mathrm{Pol}^0_{d_i}(B,\mathbb{R})$, $i = 1,\ldots,k$.

Proof. We will use induction on h; when $h = 0$ the statement is trivial. Take r_0 to be the maximum of the integers $r(l,(b_1,\ldots,b_l),s)$ in Theorem 1.9 over all integers $l \leq kw$ and all l-tuples $(b_1,\ldots,b_l)$ of nonnegative integers with $\sum_{j=1}^{l} b_j \leq w\sum_{i=1}^{k} d_i$. By induction on h, let r be the maximum of the integers $r(l,(d_1,\ldots,d_l),h-1,w,r_0)$ in the assertion of Theorem 1.10 over all integers $l \leq kw$ and all l-tuples $(b_1,\ldots,b_l)$ of nonnegative integers with $\sum_{j=1}^{l} b_j \leq w\sum_{i=1}^{k} d_i$. Let $\varphi_i \in \mathrm{GPol}^0_{d_i,h,w}([1,r],\mathbb{R})$, $i = 1,\ldots,k$. For each i reprsent φ_i in the form

$$\varphi_i = \sum_{j=1}^{m_i}[\varphi_{i,j,1}]\cdots[\varphi_{i,j,l_{i,j}}]\varphi_{i,j,0},$$

where for every i,j we have $\varphi_{i,j,0} \in \mathrm{Pol}^0_{d_{i,j,0}}([1,r],\mathbb{R})$ and for every $t \geq 1$ we have $\varphi_{i,j,t} \in \mathrm{GPol}^0_{d_{i,j,t},h-1,w}([1,r],\mathbb{R})$ with

$$\textstyle\sum_{j=1}^{m_i}(l_{i,j}+1) \leq w \text{ for all } i \text{ and } \sum_{t=0}^{l_{i,j}} d_{i,j,t} \leq d_i \text{ for all } i,j,$$

so that

$$\textstyle\sum_{i=1}^{k}\sum_{j=1}^{m_i}(l_{i,j}+1) \leq kw \text{ and } \sum_{i=1}^{k}\sum_{j=1}^{m_i}\sum_{t=0}^{l_{i,j}} d_{i,j,t} \leq w\sum_{i=1}^{k} d_i.$$

By the choice of r there exists a disjoint r_0-subcollection $B_0 \subset \mathcal{F}([1,r])$ such that $\varphi_{i,j,t\downarrow B_0} \in \mathrm{Pol}^0_{d_{i,j,t}}(B_0,\mathbb{R})$ for all i,j,t. Then by the choice of r_0 there exists a disjoint

s-subcollection B of B_0 such that for all i,j,t, $\big[\varphi_{i,j,t}\!\downarrow_B\big] \in \mathrm{Pol}^0_{d_{i,j,t}}(B,\mathbb{Z})$. Hence for every i,

$$\varphi_i\!\downarrow_B = \sum_{j=1}^{m_i} \big[\varphi_{i,j,1}\!\downarrow_B\big] \cdots \big[\varphi_{i,j,l_{i,j}}\!\downarrow_B\big] \varphi_{i,j}\!\downarrow_B \in \mathrm{Pol}^0_{d_i}(B,\mathbb{R}).$$

■

Combining Theorem 1.10 and Corollary 1.2, we obtain:

Theorem 1.11. *For any $k,d,h,w \in \mathbb{N}$ there exists $r = r(k,d,h,w) \in \mathbb{N}$ such that for any $\varphi_1,\ldots,\varphi_k \in \mathrm{GPol}^0_{d,h,w}([1,r],\mathbb{R})$ there exists a nonempty $\alpha \in \mathcal{F}([1,r])$ such that $\|\varphi_i(\alpha)\| < \varepsilon$, $i = 1,\ldots,k$.*

Let $X = G/\Gamma$ be a k-dimensional compact nilmanifold; we may and will assume that X is connected. (Any nilmanifold is a subnilmanifold of a connected one.) Let ρ be a metric on X (induced by a metric on G compatible with the Lie group structure thereon). Fix a point $x_0 \in X$, and let $\tau = (\tau_1,\ldots,\tau_k)\colon X \longrightarrow [0,1)^k$ be Maltsev's coordinates on X centered at x_0. The inverse mapping τ^{-1} is continuous, and the distance $\rho(x,x_0)$ from $x \in X$ to x_0 is continuous with respect to the distance from $\tau(x)$ to the set of vertices $\{0,1\}^k$ of the cube $[0,1]^k$. (See, for eaxmple, [BL2], Section 1.5.)

Let g be an *(l-parameter) polynomial sequence* in G, that is, a mapping $g\colon \mathbb{Z}^l \longrightarrow G$ of the form $g(n) = T_1^{p_1(n)} \cdots T_b^{p_b(n)}$, $n \in \mathbb{Z}^l$, where $T_1,\ldots,T_b \in G$, $p_1,\ldots,p_b$ are polynomials $\mathbb{Z}^l \longrightarrow \mathbb{Z}$; we define $\text{n-deg}\, g$, *the naive degree of* g, as $\max_{i=1}^b \deg p_i$. Then for each $i = 1,\ldots,k$, the sequences $\psi_i(n) = \tau_i(g(n)x_0)$, $n \in \mathbb{Z}^c$, of coordinates of x_0 under the action of g are open $[0,1)$-valued generalized polynomials, with parameters depending only on X and $\text{n-deg}\, g$ (see [BL2], Theorem A and Theorem A**), and if $g(0) = 1_G$, these polynomials can be assumed to be constant-free. For any polynomial mapping $\varphi \in \mathrm{Pol}_d([1,r]),\mathbb{Z}^c)$, the composition mappings $\psi_i \circ \varphi\colon \mathcal{F}([1,r]) \longrightarrow [0,1)$, $i = 1,\ldots,k$, are open constant-free generalized polynomial mappings, with parameters only depending on X, d, and $\text{n-deg}\, g$. From Theorem 1.11 we now obtain the following result:

Theorem 1.12. *Let $X = G/\Gamma$ be a nilmanifold with metric ρ. For any $a,d \in \mathbb{N}$ and $\varepsilon > 0$ there exists $r = r(a,d,\varepsilon) \in \mathbb{N}$ such that for any l, any l-parameter polynomial sequence g in G with $\text{n-deg}\, g \le a$ and $g(0) = 1_G$, any $x_0 \in X$, and any $\varphi \in \mathrm{Pol}^0_d([1,r],\mathbb{Z}^l)$ there exists a nonempty $\alpha \in \mathcal{F}([1,r])$ such that $\rho\big(g(\varphi(\alpha))x_0, x_0\big) < \varepsilon$.*

Remark. Theorem 1.12 easily extends to *generalized polynomial sequences* in nilpotent groups, that is, to sequences of the form $g(n) = T_1^{p_1(n)} \cdots T_b^{p_b(n)}$ where p_i are generalized polynomials $\mathbb{Z}^l \longrightarrow \mathbb{Z}$.

2. IP^*_0-recurrence implies approximability by nilsystems

In this section we prove the second statement of Theorem 0.2. Let (X,ρ) be a compact metric space, T be a self homeomorphism of X, and assume that (X,T) is IP^*_0-recurrent. Then, in particular, (X,T) is IP^*-recurrent, so by Theorem 0.1, (X,T) is distal, and thus is a disjoint union of minimal subsystems (see [F], corollary to Theorem 8.7). Hence, we may assume that (X,T) is minimal.

Now, by the way of contradiction, assume that a minimal system (X,T) is not a pre-nilsystem, that is, not an inverse limit of nilsystems; our goal is to show that there exists a point $x \in X$ and $\varepsilon > 0$ such that for every $r \in \mathbb{N}$ there exists a linear mapping $\varphi \in \mathrm{Lin}([1,r],\mathbb{Z})$ such that $\rho(T^{\varphi(\alpha))}x, x) > \varepsilon$ for every nonempty $\alpha \subseteq [1,r]$.

We will use the following result ([HKM] Theorem 1.3 and Corollary 4.2): for any r, the maximal r-step pro-nilfactor of (X,T) is defined by a closed T-invariant equivalence relation $\mathbf{RP}^{[r]} \subseteq X^2$ (called *the regionally proximal relation of order* r), with $(x_0,y_0) \in \mathbf{RP}^{[r]}$ iff for any $\delta > 0$ there exists a point $x \in X$ and a mapping $\varphi \in \mathrm{Lin}([1,r],\mathbb{Z})$ such that

$$\rho(x,x_0) < \delta \text{ and } \rho(T^{\varphi(\alpha)}x, y_0) < \delta \text{ for all nonempty } \alpha \subseteq [1,r]. \tag{2.1}$$

Our assumption that (X,T) is not a pre-nilsystem is equivalent to the assumption that $\bigcap_{r=1}^{\infty} \mathbf{RP}^{[r]} \neq \Delta$, where Δ is the diagonal of X^2. Fix $(x_0,y_0) \in \bigcap_{r=1}^{\infty} \mathbf{RP}^{[r]}$ with $x_0 \neq y_0$. Let $\varepsilon = \inf_{n\in\mathbb{Z}} \rho(T^n x_0, T^n y_0)$; since (X,T) is distal, we have $\varepsilon > 0$. Since (X,T) is minimal, the orbit $\{T^n x_0\}_{n\in\mathbb{Z}}$ of x_0 is dense in X. Let $r \in \mathbb{N}$ and let $U \subseteq X$ be an open set. Choose $n \in \mathbb{Z}$ such that $T^n x_0 \in U$ and choose $\delta > 0$ such that $\rho(T^n x, T^n y) < \varepsilon/3$ whenever $\rho(x,y) < \delta$. Find $x \in X$ such that (2.1) holds and $T^n x \in U$. Then $\rho(T^n x, T^n x_0) < \varepsilon/3$ and $\rho(T^{\varphi(\alpha)} T^n x, T^n y_0) < \varepsilon/3$ for all nonempty $\alpha \subseteq [1,r]$, and since $\rho(T^n x_0, T^n y_0) \geq \varepsilon$, we have that $\rho(T^{\varphi(\alpha)} T^n x, T^n x) > \varepsilon/3$ for all nonempty $\alpha \subseteq [1,r]$. This proves that for any $r \in \mathbb{N}$ the open set

$$R_r = \big\{x \in X : \text{there exists } \varphi \in \mathrm{Lin}([1,r],\mathbb{Z}) \text{ such that } \rho(T^{\varphi(\alpha)}x, x) > \varepsilon/3 \text{ for all nonempty } \alpha \in [1,r]\big\}$$

is dense in X. By Baire category theorem $\bigcap_{r=1}^{\infty} R_r$ is nonempty, which gives us what we wanted – a point $x \in X$ such that for every $r \in \mathbb{N}$ there exists a mapping $\varphi \in \mathrm{Lin}([1,r],\mathbb{Z})$ such that $\rho(T^{\varphi(\alpha))}x, x) > \varepsilon/3$ for every nonempty $\alpha \subseteq [1,r]$.

Bibliography

[AGH] L. Auslander, L. Green and F. Hahn, *Flows on homogeneous spaces*, Annals of Math. Studies, vol. 53, Princeton Univ. Press, 1963.

[B] V. Bergelson, Ultrafilters, IP sets, Dynamics, and Combinatorial Number Theory, *Contemp. Math.* **530** (2010), 23-47.

[BL1] V. Bergelson and A. Leibman, Set-polynomials and polynomial extension of the Hales-Jewett Theorem, *Ann. of Math.* **150** (1999), no. 1, 33-75.

[BL2] V. Bergelson and A. Leibman, Distribution of values of bounded generalized polynomials, *Acta Math.* **198** (2007), 155-230.

[BL3] V. Bergelson and A. Leibman, Sets of large values of correlation functions for polynomial cubic configurations, preprint.

[F] H. Furstenberg, *Recurrence in Ergodic Theory and Combinatorial Number Theory*, Princeton Univ. Press, 1981.

[HKM] B. Host, B. Kra, and A. Maass, Nilsequences and a structure theorem for topological dynamical systems, *Adv. in Math.* **224** (2010), no. 1, 103-129.

[Ke1] H. B. Keynes, Topological dynamics in coset transformation groups, *Bull. Amer. Math. Soc.* **72** (1966), 1033-1035.

[Ke2] H. B. Keynes, A study of the proximal relation in coset transformation groups, *Trans. Amer. Math. Soc.* **128** (1967), 389-402.

[P] R. Pavlov, Some counterexamples in topological dynamics, *Ergod. Th. and Dynam. Sys.* **28** (2008), no. 4, 1291-1322.